Socialstyrelsen
Hälso- och sjukvårdsavdelningen

Janne Larsson
Snöbollsgränd 22
129 45 Hägersten

Angående uppgift ur handlingen Förslag till nationellt program för suicidprevention

Socialstyrelsen har i e-post skickad den 16 december 2006 fått en fråga om en uppgift i Förslag till nationellt program för suicidprevention. Ni vill ha en referens till följande uppgifter:

”Av de personer med depressionsdiagnos som suiciderar är mer än 80 procent obehandlade vid tidpunkten för dödsfallet. Personer med depression är ofta obehandlade eller underbehandlade även efter ett suicidförsök.”
I en artikel av Isacsson G, Bergman U och Rich CL (1996) uppskattas det årliga antalet suicid med depression som tagit antidepressiva medel till 120 av 1 000 fall i Sverige (12%) och antalet årliga suicid som inte tar antidepressiva medel till 880 av 1 000 fall (88%). Detta stöds också av internationella studier. Enligt Goodwin & Jamison (1999) har 15-19 procent av patienter som suiciderat tidigare sjukhusvårdats för depression. Mann et al (2005) påpekar i en review att depression i allmänhet är obehandlad eller underbehandlad även efter ett suicidförsök. Uppgiften stöds av en studie av Oquendo, Kamali & Ellis 2002.

Referenser:

Isacsson G, Bergman U & Rich CL. Epidemiological data suggest antidepressants reduce suicide risk among depressives. Journal of Affective Disorders 1996;41:1-8.

Mann et al. Suicide Prevention Strategies. A Systematic Review. JAMA 2005;294(16): 2064-74.

Hirschfeld RMA, Keller M, Panico S, et al. The National Depressive and Manic-Depressive Association consensus statement on the undertreatment of depression. JAMA 1997;277:333-340.

Goodwin FK, Jamison KR. Manic-depressive illness. New York: Oxford University Press 1999.

Coyle JT, Pine DS, Charney DS, et al. Depression and bipolar support alliance consensus statement on the unmet needs and treatment of mood disorders in children and adolescents. J Am Acad Child Adolesc Psychioatry 2003;42:1494-1503.

Oquendo MA, Kamali M, Ellis SP, et al. Adequacy of antidepressant treatment after discharge and the occurrence of suicidal acts in major depression: a prospective study. Am J Psychiatry 2002;159:1746-1751.

Med vänlig hälsning

Anders Hedberg